



Open Access

INVITED REVIEW

Androgens and estrogens in skeletal sexual dimorphism

Michaël Laurent^{1,2,3}, Leen Antonio^{1,4,5}, Mieke Sinnesael⁴, Vanessa Dubois¹, Evelien Gielen^{2,3,6}, Frank Classens¹, Dirk Vanderschueren^{4,5,6,7}

Bone is an endocrine tissue expressing androgen and estrogen receptors as well as steroid metabolizing enzymes. The bioactivity of circulating sex steroids is modulated by sex hormone-binding globulin and local conversion in bone tissue, for example, from testosterone (T) to estradiol (E2) by aromatase, or to dihydrotestosterone by 5 α -reductase enzymes. Our understanding of the structural basis for gender differences in bone strength has advanced considerably over recent years due to increasing use of (high resolution) peripheral computed tomography. These microarchitectural insights form the basis to understand sex steroid influences on male peak bone mass and turnover in cortical vs trabecular bone. Recent studies using Cre/LoxP technology have further refined our mechanistic insights from global knockout mice into the direct contributions of sex steroids and their respective nuclear receptors in osteoblasts, osteoclasts, osteocytes, and other cells to male osteoporosis. At the same time, these studies have reinforced the notion that androgen and estrogen deficiency have both direct and pleiotropic effects via interaction with, for example, insulin-like growth factor 1, inflammation, oxidative stress, central nervous system control of bone metabolism, adaptation to mechanical loading, etc., This review will summarize recent advances on these issues in the field of sex steroid actions in male bone homeostasis.

Asian Journal of Andrology (2014) 16, 213–222; doi: 10.4103/1008-682X.122356; published online: 20 January 2014

Keywords: androgen receptor; bone mineral density; estradiol; estrogen receptor; male; osteoblast; osteoclast; osteocyte; osteoporosis; sex hormone-binding globulin; testosterone

INTRODUCTION

Bone is an endocrine tissue sensitive to androgens and estrogens.¹ The epidemiology and clinical approach to male osteoporosis have recently been reviewed elsewhere.² Here, we will review the literature on the role of sex steroids in male bone health and more specifically in osteoporosis, the most common metabolic bone disease. We will focus on human data with support from recent evidence from knockout mouse models. The latter are covered in more detail elsewhere in this issue (See Rana *et al.* in this theme issue).

Men contribute considerably to the disease burden of osteoporosis

Although still perceived by the public, patients and most physicians as a typically female condition, men account for a substantial proportion of the burden of osteoporosis. At age 50, the remaining lifetime risk of osteoporotic fractures is 20%–25% in Caucasian men versus 45%–55% in women.² About one-third of the 9 million osteoporotic fractures worldwide occurs in men.³ For hip fractures, a review of worldwide epidemiological data recently showed that while incidence varied more than 10-fold between countries, men almost invariably have twofold lower risk (except perhaps in regions where incidence is very low).⁴ Male osteoporosis is responsible for about 25% of the economic burden of osteoporosis in the United States (US\$17 billion total costs) and Canada (2.3 billion Canadian dollars).^{5–7} Nevertheless, osteoporosis remains even more underdiagnosed and undertreated in men compared to women, regardless of whether it is evidenced by low bone mineral

density (BMD),⁸ a low-energy fracture^{9,10} or absolute 10-year fracture probability.^{11,12}

Basic sex steroid physiology in relation to bone health

Metabolism and regulation of circulating sex steroids

Sex steroid serum concentrations are determined by their synthesis in the gonads and adrenals as well as by catabolic enzyme activity. Testosterone (T), the principal circulating androgen, can be aromatized to estradiol (E2, the principal estrogen) or 5 α -reduced to dihydrotestosterone (DHT), a non-aromatizable androgen. Thus, in bone T may stimulate the androgen receptor (AR), either directly or as DHT, as well as estrogen receptors (ER) following aromatization.¹³ The synthesis of sex steroids is under hypothalamic-pituitary feedback control via gonadotropins (follicle stimulating hormone (FSH) and luteinizing hormone (LH)). Several high-profile reports have suggested that FSH contributes to hypogonadal bone loss in female rodents,^{14,15} but other groups have failed to confirm the presence of FSH receptors in bone cells or a physiological effect thereof.^{16–19} Gonadotropin inhibition also does not seem to influence bone turnover markers independent of sex steroid levels in men.²⁰

Serum bioactivity of sex steroids is restricted in humans by their high affinity binding to sex hormone-binding globulin (SHBG). About 45% of T, the principal circulating androgen, is bound to SHBG in normal men; only the non-SHBG-bound fraction is considered

¹Laboratory of Molecular Endocrinology, Department of Cellular and Molecular Medicine; ²Gerontology and Geriatrics, Department of Clinical and Experimental Medicine, KU Leuven, Leuven, Belgium; ³Geriatric Medicine, University Hospitals Leuven, Leuven, Belgium; ⁴Clinical and Experimental Endocrinology, Department of Clinical and Experimental Medicine, KU Leuven, Leuven, Belgium; ⁵Endocrinology and Andrology; ⁶Centre for Metabolic Bone Diseases; ⁷Laboratory Medicine, University Hospitals Leuven, Leuven, Belgium.

Correspondence: Dr. M. Laurent (michael.laurent@med.kuleuven.be)

Received: 06 June 2013; Accepted: 02 July 2013

bioactive. Most of this fraction is bound nonspecifically to albumin and other carrier proteins, while only about 2% of T and E2 and <1% of DHT circulates freely in normal men.²¹ According to the free hormone hypothesis however, this constitutes the most physiologically important fraction.²² Others have pointed to the limitations of this theory and have pleaded to “stop looking backwards” and focusing exclusively on free sex steroid levels. Indeed, even if only a small portion of the protein-bound sex steroids is physiologically active, it is likely to have more influence than the 1%–2% that is free.²³

Sex steroid nuclear receptor signaling

Sex steroids classically act via their nuclear receptors (NRs: AR, ER α and ER β), which bind directly or indirectly via other proteins to cognate DNA response elements (androgen response elements and estrogen response elements), recruit coactivators and other regulatory proteins, and influence gene transcription.²⁴ Two types of androgen response elements exist (classical and selective), but mice with abrogated AR binding to selective androgen response elements only show a sex organ phenotype.²⁵ Nonclassical signaling pathways of these NRs have also been described but remain somewhat controversial (especially for AR). These have not yet been successfully translated (therapeutically) to studies in humans, and will therefore not be further discussed here.

Androgen and estrogen deficiency also have important indirect effects via other signaling pathways in bone (e.g. insulin-like growth factor I (IGF-1)²⁶), potentially via AR and ER in other tissues like muscle, fat, and the nervous system, or via, for example, decreased physical activity in AR knockout (ARKO) mice.^{27,28} Therefore, caution needs to be exerted regarding the roles of AR and ER specifically in bone cells when interpreting the results of gonadectomy or ARKO/ERKO models. These limitations have been overcome recently using Cre/LoxP mouse models (see below). The fact that these cell-specific models show less dramatic bone phenotypes than their global ARKO/ERKO counterparts confirms the importance of pleiotropic actions of sex steroids on bone.

STRUCTURAL BASIS OF MALE BONE STRENGTH

The musculoskeletal system is sexually dimorphic, being on average (but obviously not in all cases) larger and more robust in men compared to women. Bone strength is determined by peak bone mass (PBM) acquisition in young adulthood and subsequent turnover in different compartments (cortical and trabecular bone), resulting in gender differences in bone length, bone mineral density (BMD), geometry, and microarchitecture. For other determinants like material properties, gender differences require further investigation. Osteoporotic fracture risk is not only determined by purely skeletal factors but also by risk of falls, which should also be evaluated and treated in osteoporotic men.^{29,30}

In the following paragraphs, we will discuss the structural basis of male bone strength. It is well-known that men have higher peak areal BMD (aBMD) in young adulthood and aBMD declines slower as they age.³¹ However, this should be interpreted with caution because of the projectional nature of dual energy X-ray absorptiometry (DXA), which leads to potential confounding by bone size, geometry, and porosity rather than true volumetric density (vBMD). In this respect, quantitative computed tomography (qCT), especially high resolution peripheral qCT (HR-pqCT), has recently led to a major advance in our understanding of the gender dimorphism in the structural determinants of bone strengths.

Bone size

Men are on average taller due to the actions of sex steroids on the growth plate (see below). From a biomechanical point of view, longer

bones (which are also wider and have greater bone mass) are more resistant to bending. However, several cohort studies have found a positive association between height and risk of low-energy fractures, although the results in men are scant and only bordered significance.^{32,33} This may be explained by greater loads when falling from a larger height, longer arms of moment, as well as by relatively thinner cortices and greater cortical porosity in larger bones; although definitive data in men are currently lacking.³⁴

Peak bone mass

Bone mineral mass can be low in old age because insufficient PBM attainment during skeletal growth (modeling phase) or due to an imbalance in the normally coupled process of bone turnover (remodeling phase), either by excessive bone resorption or decreased bone formation. The risk of many chronic diseases which manifest in old age, including osteoporosis, is determined at least in part by childhood or even prenatal experiences. PBM is largely (60%–80%) genetically determined, but the remainder is determined by environmental factors, some of which (e.g. maternal age and vitamin D status during pregnancy) are potentially amenable to interventions.^{35,36} Theoretically, small increases in PBM translate into a large delay in osteoporosis onset,³⁷ although it remains to be established whether bone mass increments early in life can be sustained into old age.³⁸

The age of PBM or peak bone density differs between studies and depends on the measured site. In a prospective study with 15 years of follow-up, BMC in boys plateaued first at the femoral neck, followed by the total hip, lumbar spine, and whole body, respectively at the age of 2, 3, 4, and 6 years after peak height velocity (PHV, average age 14 in men).³⁹ Hip aBMD decreases early after it peaks.⁴⁰ Longitudinal studies across a wider age range have however estimated that lumbar spine BMD may peak as late as age 33 and 40 in men and women, respectively. Bone area increments are followed 1–2 years later by mineralization, and there are little differences between men and women when the age of PHV is used as the reference.⁴¹ The peripubertal years (age 12–16) contributed 33%–46% to adult-level BMC, demonstrating the importance of this window of opportunity for interventions to optimize PBM.³⁹ There is considerable variation in the timing of male puberty, but neither early nor late normal puberty seem to compromise PBM.⁴²

Bone geometry and microarchitecture

Men develop wider bones even after adjustment for height because of greater periosteal apposition in the appendicular skeleton, whereas girls predominantly decrease their endosteal perimeter.¹ As a result, cortical bone in men is placed further away from the neutral axis, providing greater resistance to bending, but no difference in size-independent biomechanical indices.^{43,44} Bone width predicts fractures in older men independently of aBMD, and men with low bone width combined with BMD T-scores <–1 have similar fracture incidence as men with T-scores <–2.⁴⁵

Recently, most cross-sectional studies using HR-pqCT have documented cortical and trabecular bone across the lifespan in more detail (**Figure 1**).

At PBM, men have higher trabecular bone volume fraction (**Figure 1a**) due to thicker (**Figure 1c**), more plate-like trabeculae. On the other hand, cortical porosity is greater (not shown) and cortical vBMD slightly lower (**Figure 1f**), while trabecular vBMD (not shown) and cortical thickness (**Figure 1e**) are not yet greatly different.^{43,46–48} Increasing cross-sectional area and cortical thickness, ongoing cortical mineralization and decreasing porosity contribute to ongoing aBMD and estimated strength increases during




Figure 1: Structural determinants of bone strength in men based on high resolution peripheral computed tomography (HR-pqCT) at the ultradistal radius. Adapted, with permission, from Khosla *et al.*⁴⁶ Tb.Th: trabecular thickness; BV/TV: trabecular bone volume/tissue volume; Tb.N: trabecular number; Tb.Sp: trabecular separation; Cort vBMD: cortical volumetric bone mineral density; Ct.Th: cortical thickness.

the 3rd decade of life in men, whereas in women cortical perimeter and strength do not increase during young adulthood.^{40,49}

Both cortical and trabecular vBMD losses are more pronounced with age in women, although rates of trabecular bone loss vary across sites.^{43,50} With ageing, endocortical resorption outpaces periosteal apposition leading to a thinner cortex in both sexes, but older women show dramatically greater medullary expansion and thus cortical thinning (Figures 1e and 3).^{43,51–53} This was nicely illustrated in one prospective pqCT study (Figure 2), which also demonstrated the limitations of capturing bone geometrical changes across the lifespan from cross-sectional data.⁵¹

Cortical porosity is greater in young men, but increases faster in women, especially after menopause, resulting in similar porosity after age 50.^{43,44} Cortical vBMD decreases from midlife in women, but only after age 75 in men, whereas trabecular bone loss starts almost immediately after PBM in both sexes (Figure 1a) but more so in women, with acceleration during perimenopause.^{50,53} Bone loss in men is more due to decreased formation rather than increased resorption, therefore trabeculae become thinner (Figure 1c) but less perforated, disconnected and thus widely spaced (Figure 1b and 1d) compared to postmenopausal women.^{46,54} Poor trabecular microarchitecture predisposes men to multiple and severe vertebral and peripheral fractures.⁵⁵ At the femoral neck, men have higher aBMD despite lower vBMD, due to greater length and femoral neck area, showing




Figure 2: Schematic model of bone geometry changes at the tibia during adult lifespan in men and women. When comparing young women and men, men have greater cortical thickness due to greater periosteal perimeter (shown in black), while women mainly decrease their endocortical perimeter (also shown in black). With ageing, cortical thinning results from endocortical bone resorption in both genders (difference compared to age 20 shown in blue), but this is much more pronounced in women, and ongoing periosteal apposition (difference compared to age 20 shown in red) in both genders is unable to compensate for the differences. On the other hand, bone is placed further away from the central axis in men at all ages, and this dramatically improves biomechanics. Adapted, with permission, based on data from Lauretani *et al.*⁵¹

the importance of geometry over mineralization in the decreased hip fracture risk in men.⁵⁶ This also explains why men and women have the same strength for the same femoral neck aBMD: because of the offsetting effects of higher bone area and lower vBMD in men.⁵⁷

In conclusion, detailed imaging techniques have greatly improved our understanding of the structural basis for gender differences in bone strength. How and to what extent sex steroids regulate this sexual dimorphism in cortical and trabecular bone will be discussed below.

SEX STEROID SIGNALING IS KEY TO MUSCULOSKELETAL SEXUAL DIMORPHISM

Are gender differences in bone phenotype due to sex steroid signaling or other pathways? Sex steroids and especially estrogens were once considered a central hub on which bone signaling pathways converged in both genders,⁵⁸ but this unitary estrogen-centric view has been modified.^{59,60} Both at PBM and during ageing, BMD, bone geometry, bone turnover, and skeletal muscle mass in men have now been associated with multiple hormones (e.g., IGF-1 and IGF binding proteins,^{61–64} endogenous PTH,^{63,65} vitamin D,^{66,67} and thyroid hormone within the normal range),^{68,69} immunological, and other pathways in bone (e.g., CRP,⁷⁰ RANK/RANKL/OPG), oxidative stress,^{71–73} and classical ageing pathways,⁶⁰ *etc.* Mostly similar associations have however been reported in women, and a recent meta-analysis of genome-wide BMD association studies failed to identify any significant gene-by-sex interactions.⁷⁴ Fundamental gender differences in key bone signaling pathways therefore seem unlikely.

Conversely, the musculoskeletal system of transsexual men shows a dramatic shift from the female to male phenotype after ovariectomy and prolonged T treatment.⁷⁵ Similarly, XY women with complete androgen insensitivity syndrome due to inactivating AR mutations have reduced BMD and a bone geometry intermediate between male and female, and estrogen treatment does not induce periosteal bone apposition in these subjects.⁷⁶ BMD is however much lower in both XY and XX women with gonadal dysgenesis, implying that gonadal status or sex steroids are more important than chromosomal determinants.⁷⁷ Overall, we can conclude that androgens (or AR mediated androgen action) are necessary for musculoskeletal sexual dimorphism in development and ageing, although they probably have important indirect actions on bone via

aromatization, oxidative stress,⁷⁸ proinflammatory cytokines,^{79,80} growth factors (e.g. transforming growth factor (TGF)- β , IGF-1),^{1,26,81} *etc.*

SEX STEROID REGULATION OF MALE BONE METABOLISM

Sex steroids control longitudinal bone size at the growth plate

It is generally accepted that delayed estrogen-mediated closure of epiphyseal growth plate cartilage contributes to greater bone length in men. T on the other hand probably also stimulates height velocity mainly via aromatization and estrogen-mediated pituitary growth hormone release. Non-aromatizable androgens increase growth rate in boys without altering serum GH/IGF-1, possibly via the AR in chondrocytes and local IGF-1 signaling in the growth plate.^{1,82} Pubertal height velocity acceleration and subsequent growth plate closure seem to be absent in men with inactivating ER α mutations⁸³ or aromatase deficiency^{84–87} who show steady, continuous growth, suggesting that estrogens have a dominant role in these processes. A recent study using ER α KO mice with no residual truncated isoforms confirmed the continuing longitudinal growth seen in the ER α KO (see next paragraph).⁸⁸ The fact that 46, XY girls with complete androgen insensitivity syndrome have a growth spurt similar in timing and amplitude as 46, XX girls⁸⁹ is probably due to estrogen effects, but it does not exclude a role for AR itself. Male ARKO mice have been anecdotally noted to have increased femur length,⁹⁰ but in men the effects of increased estrogen due to aromatization probably override any possible direct effect of T on the growth plate. On the other hand, further research on androgen regulation of (growth plate) chondrocytes would be of interest.

Estrogen deficiency: the primary mediator of bone loss in older hypogonadal men

The primary role of estrogens in postmenopausal osteoporosis was proposed by Albright *et al.*, in 1941.⁹¹ Similarly, hypogonadal men with low T were noted to have increased risk of osteoporosis and fractures.^{92,93} The historic dichotomous view that estrogens were important for bone health in women and androgens in men was however challenged by a unique report of a man with an ER α mutation.⁸³ He presented with tall stature, incomplete epiphyseal closure in adulthood and markedly decreased bone density. In 1998, investigators from the Mayo Clinic proposed a unitary model for the pathogenesis of osteoporosis with a central responsibility for estrogens, not only in post-menopausal but also in male osteoporosis and even in the negative calcium balance and secondary hyperparathyroidism in age-related osteoporosis.⁵⁸ This was based on seminal observations,⁹⁴ subsequently confirmed in multiple prospective cohort studies, of a strong association between (calculated free or bioavailable) E2, bone turnover, and bone density in community-dwelling men; whereas, the same associations with T were variable, weak, absent, or disappeared after correcting for E2 or other variables.^{95–105} On the other hand, threshold effects (i.e., greater decreases in serum E2 in women compared to relatively normal androgen levels in ageing men) probably limit the conclusions that can be drawn from such cohort studies. Indeed, in several studies, E2 (and SHBG, see below) have been associated with bone loss below a certain threshold,^{96,101,106} while this has not been confirmed in younger cohorts of men.^{105,107} In the European Male Ageing Study (EMAS) of men aged 40–79 years, the prevalence of T between 11 and 8 nmol l⁻¹ was 12.9%, while only 4.1% had T <8 nmol/l, showing that most hypogonadism is mild in cohort studies.¹⁰⁸ Nevertheless, several complimentary lines of evidence (Table 1)^{84–87,109–120} confirm that estrogens are crucial to restrain bone turnover in ageing men. Thus, the finding that T blocks orchidectomy-induced bone loss in ER α KO mice¹²¹ has been contradicted by human experiments,¹²⁰ which is possibly due to limitations of the ER α KO mouse model (which

shows high bone mass due to increased T, which also contrasts with the situation in an ER α KO man).⁸³

Low testosterone and high sex hormone-binding globulin pose additional risks

The overall conclusion thus appears to be that estrogens are more important than androgens in maintaining bone health in ageing men. Yet low T and high SHBG may still harbor additional detriments. Free or bioavailable T has been associated with BMD (at predominantly cortical sites), bone area, muscle area and strength, reduced fat mass and hip, vertebral and non-vertebral fractures in different large studies in older men.^{107,122–124} Late-onset hypogonadism (LOH) as defined by the presence of three sexual symptoms according to criteria from the EMAS appears to correlate more strongly than low T alone with low ultrasound-estimated BMD.¹²⁵ Similarly, risk for hip fractures has been found highest in men with both low E2 and T.¹²⁶ We have previously shown in mice that optimal effects of T require a functional AR.¹²⁷ These data are in agreement with findings from a man with concomitant aromatase deficiency and low T, in whom T and E2 replacement showed additive effects.¹²⁸

In the largest study, low E2 and high SHBG were independent predictors of BMD losses and fracture risk in men, but risks were highest when both bioavailable E2 and T were combined with high SHBG.^{129,130} SHBG has been associated independently of bioavailable T and E2 with improved cortical bone development in young men, but accelerated bone losses with ageing and consequently increased fracture risk.^{106,131–134} The mechanisms by which SHBG may influence bone in this apparently paradoxical manner require further investigation.¹³⁵

The role of catabolic enzymes

Not only synthetic but also catabolic enzymes control steady state sex steroid serum levels. In the Swedish MrOS study, specific androgen metabolites correlated with male BMD, while T itself did not.¹⁰² Polymorphisms in the enzymes catechol-O-methyl-transferase (COMT, an estrogen degrading enzyme) and uridine diphosphate glucuronosyltransferase 2B7 (which inactivates mainly androgens

Table 1: Evidence supporting the primary role of estrogens in bone loss in older men

Animal models	E2 is a more effective bone-sparing agent than DHT in aged male rats ¹⁰⁹
Human genetic evidence	Polymorphisms in ER (but not AR) signaling negatively influence male bone ^{110–112} Longer CAG repeats diminish AR function but may be compensated by increased T levels, paradoxically stimulating bone via aromatization ¹¹³ Men with aromatase deficiency have high T, but suboptimal bone mass which improves with estrogen (but not androgen) therapy ^{84–87}
Human observational studies	Several large cohort studies confirm relationship between E2 and male bone metabolism, while the same relationship with T is weak, absent, or disappears after correcting for E2 (but not vice versa) Increased bone turnover and fracture risk in older men receiving androgen deprivation therapy for prostate cancer may be due to a loss of substrate for aromatization, since AR antagonists alone are not detrimental for bone ^{115–117}
Human experiments	Aromatase inhibition in older men increases T, but decreases E2 and BMD ¹¹⁸ Gonadotropin inhibition and T or E2 add-back shows that E2 has the dominant anti-resorptive effect ¹¹⁹ T replacement with or without aromatase inhibition demonstrates that T alone cannot overcome male bone resorption in the absence of E2 ¹²⁰

BMD: Bone Mineral Density; CAG: Cytosine adenine guanine; DHT: Dihydrotestosterone

but also some estrogens) have been associated with increased sex steroid levels and bone geometry in young men.¹³⁶⁻¹³⁹ In another large population-based study, the COMT polymorphism was independently associated with fracture risk but not BMD, and only in men.¹⁴⁰ The activity of these and other steroid metabolizing enzymes, and whether they exert their influence mainly systemically or locally in bone, merits further study.

The contribution of 5 α -reductase type 1

Only a small percentage of DHT enters the circulation, and this fraction does not correlate with male BMD.¹⁰² 5 α -reductase (encoded by *Srd5a*) type 2 is primarily expressed in male reproductive tissues including the prostate; whereas, type 1 and 3 are expressed in many tissues including reproductive tissues and bone.¹⁴¹ The 5 α -reductase inhibitor finasteride inhibits type 2 and 3; whereas, dutasteride inhibits all three. It is therefore not unexpected that finasteride does not affect bone density, turnover, fracture risk, or muscle anabolic effects of androgens.¹⁴²⁻¹⁴⁸ In contrast, male *Srd5a1*^{-/-} mice have recently been characterized as having reduced bone mass and muscle strength despite normal androgen serum levels.¹⁴⁹ Human RCTs however have not shown effects of dutasteride on bone metabolism,^{145,150} nor does dutasteride influence the anabolic effects of T on muscle.¹⁵¹ Whether or not 5 α -reductase type 1 activity contributes to male bone mass, thus merits further investigation.

Meanwhile, aromatizable androgens may be used preferentially to simultaneously compensate androgen and estrogen deficiency in men with LOH. In a small randomized controlled trial (RCT), DHT gel treatment in older men with low T suppressed endogenous T without affecting E2 or BTMs,¹⁵² while another RCT showed decreased E2 as well as spinal BMD.¹⁵³ Therefore, the main effect of androgens or selective AR modulators (SARMs) may depend on their suppression of estrogens. We have previously shown in rodents that androgens and estrogens apparently act via different pathways and that their combination might be beneficial compared to each strategy alone,^{154,155} although this requires confirmation in men with LOH.

RELATIVE CONTRIBUTIONS OF ANDROGEN RECEPTOR, ESTROGEN RECEPTOR α , AND ESTROGEN RECEPTOR β IN CORTICAL VS TRABECULAR BONE

Although observational studies in humans are important to determine the influence of sex steroids on male bone, the study of the respective contributions of AR and ER require knockout animal models as well as confirmation in rare human genetic disorders. These studies have revealed unexpected complexity regarding the roles of these NRs in different bone compartments.²⁶

Both AR and ER α are required for optimal periosteal bone expansion,^{90,156} although estrogens limit periosteal expansion in early puberty, probably because of time-specific and independent effects on IGF-1.^{26,81} For trabecular bone development on the other hand, AR is solely responsible.^{90,156} Indeed, ARKO decreases cancellous bone, while systemic ER α KO increases it (probably due to increased androgen levels in this animal model)^{121,157} and combined AR/ER α KO does not decrease trabecular bone more than ARKO alone.⁹⁰ Compared to wild-type females, male pubertal ARKO mice have equal length, decreased trabecular bone, and identical cortical bone parameters; showing that androgens are required for optimal bone development, especially trabecular but also cortical, but not for longitudinal growth.¹⁵⁸

These results are confirmed in humans: estrogen therapy in young aromatase-deficient men indeed improves cortical thickness and area; however, without increasing trabecular vBMD.⁸⁶ Nevertheless, more recent animal studies using cell-specific ER α KO models do also suggest a role for ER α in trabecular bone formation (see below). ER β plays a role in female bone health,^{157,159,160} but male ER β KO mice have normal bones and ER α β KO show no difference to ER α KO alone.¹⁵⁴

ACTION MECHANISMS OF AR AND ER α IN SPECIFIC BONE CELLS

Androgens control osteoblasts and osteocytes, while estrogens also regulate osteoclasts

Cell culture experiments from ARKO mice have suggested that AR controls mainly osteoblasts and their indirect control of osteoclastogenesis.¹⁶¹ However, as mentioned above, global ARKO/ERKO models or gonadectomy/antagonist studies preclude definitive conclusions about the cellular targets of AR and ER in male bone. Indirect evidence suggested that AR and ER signaling targets mainly osteoblasts and osteoclasts, respectively. These assumptions have only recently been verified using Cre/LoxP mouse models, although this requires careful verification that neither the Cre driver nor the floxed genotypes have effects themselves (see Rana *et al.* in this theme issue).

Androgen receptor in the osteoblast lineage

Osteocalcin-Cre driven ARKO revealed that androgens stimulate mineralizing osteoblasts, thus indirectly inhibiting cortical and trabecular bone resorption, especially during times of bone accrual.¹⁶² Col2.3-Cre driven ARKO showed that mature osteoblasts contribute to trabecular bone maintenance.¹⁶³ However, periosteal apposition was not influenced by AR deficiency in these mature cells, probably because the periosteum contains more pre- and proliferating osteoblasts. Indeed, AR overexpression in immature osteoblasts increases periosteal and decreases endosteal bone formation; whereas, osteogenesis was inhibited at both envelopes following AR overexpression in mature osteoblasts.^{164,165} Recently, we have shown in Dmp1-Cre mice that loss of AR in osteocytes results in similar, moderate impairment of trabecular bone maintenance.¹⁶⁶ Importantly, AR expression increased with the maturation of osteoblasts towards osteocytes.¹⁶⁶ These different phenotypes suggest that AR has a direct role across the entire osteoblast lineage.

Androgen receptor and estrogen receptor alpha in osteoclasts

Clearly, AR signaling has important indirect restraining effects on osteoclasts, for example by controlling cytokine production in bone marrow stromal cells.⁷⁹ AR has been detected *in vitro* and with immunohistochemistry in rodent osteoclasts albeit at very low levels, and it may be absent in human osteoclasts.^{1,167,168} Several though not all studies have suggested that androgens also directly suppress *in vitro* osteoclast formation from hematopoietic precursors.^{1,168-171} Although they could not confirm the presence of AR in osteoclasts, the group of Kato found to their surprise that cathepsin K-Cre-driven ARKO also induces bone resorption (although this data has only been published in abstract),¹⁷² in a similar way as they and others have reported for osteoclast-specific ER α KO in female (but not male) mice.^{173,174} Thus, a direct *in vivo* role for AR in osteoclasts merits independent confirmation.

Estrogen receptor alpha in osteoblasts

ER α has recently been established to have a role in osteoblasts and osteocytes too. It was already known that osteoblast-specific overexpression of aromatase increases bone mass in male mice.¹⁷⁵ Prx1

Cre and Osterix-Cre mice have now been used to selectively excise ER α from pluripotent mesenchymal progenitors in the limb bud of the appendicular skeleton, and respectively, osteoblast progenitors. These mice showed mainly cortical bone deficits resulting from decreased periosteal bone formation, although cortical bone deficits were overcome during adulthood in Prx1-Cre ER α KO males.¹⁷⁶ Deletion of ER α using the Col1a1 deleter did not affect cortical or trabecular bone. However, this should not be taken as evidence that ER α has no role downstream in osteoblast differentiation. Indeed, osteocalcin-Cre ER α KO decreased trabecular bone in males, and both trabecular and cortical bone in females.¹⁷⁷ Dmp1-Cre ER α KO males also showed decreased bone formation and less trabecular bone, but there was no effect on cortical bone, or any effect in females. The authors concluded that the trabecular bone-sparing effects of estrogens are mediated by osteocyte ER α in males, but probably by osteoclast ER α in females.¹⁷⁸ Interestingly, both ER α KO¹⁷⁹ and ARKO¹⁸⁰ have been known to influence the response to mechanical loading, but the response to mechanical loading was unaltered in the respective osteocyte-specific knockout mice.^{166,178}

In summary, these studies in male mice suggest that both AR and ER α are required for optimal cortical bone expansion via actions in immature osteoblasts, and trabecular bone maintenance via actions in more differentiated osteoblasts and osteocytes (Figure 3).

Indirect effects of androgen receptor and estrogen receptors on bone via muscle, fat, and the nervous system

The interaction of bone with muscle, adipose, and neural systems is increasingly studied. Tissue-specific ARKO models are discussed in more detail in the accompanying review by Rana *et al.* in this theme issue.

Bone-muscle interaction

Male ARKO mice have impaired muscle development,¹⁸¹ and additional ER α KO further reduces muscle mass.⁹⁰ However, muscle-specific ARKO mice did not show altered bone metabolism and only slightly reduced peripheral skeletal muscle mass, possibly because only perineal muscles display high AR content and androgen regulation in mice.^{182,183} This contrasts with the well-known anabolic effects of androgens on human muscle.¹⁸⁴ Thus, although the effects of androgens on bone in mice are unlikely to be mediated via the AR in muscle, it remains unclear whether sarcopenia in older men with LOH contributes directly or is merely associated with bone loss and fractures.^{2,185,186}

Bone-fat interaction

Clinical evidence suggests a positive association between bone and fat, but mainly in females, possibly because adipocyte aromatase activity influences circulating E2 or because of increased gravitational loading.^{2,185} Fat mass is principally regulated by estrogens since double knockouts have similar adiposity compared to ER α KO alone,⁹⁰ although androgens also have clear lipolytic activity, and male ARKO mice have increased adiposity.²⁸ A link between glucose, insulin, and bone metabolism has also been suggested in mice, and male bone metabolism is altered in diabetes and the metabolic syndrome. Thus, whether AR or ER signaling in adipocytes modulates sex steroid effects on bone would be of interest.

Central nervous system control of bone mass

Central nervous system regulation of bone mass has been demonstrated recently with often opposite effects to peripheral signaling. This also seems to be the case for estrogens, since neuron-specific ER α KO using nestin-Cre increases bone formation via leptin.¹⁸⁷ Conditional inactivation of AR in the nervous system of mice however disrupts the somatotrophic axis as evidenced by growth retardation and twofold lower serum IGF-1, without relative differences in total bone mass or body composition.¹⁸⁸ This reiterates that the skeletal sexual dimorphism resulting from sex steroids is dependent on indirect effects via growth hormone, IGF-1 and IGF binding proteins in brain, bone, and liver.^{26,81,189}

CONCLUSIONS

The musculoskeletal system is more robust in men, and sex steroid signaling remains essential for this sexual dimorphism. The advent of high resolution imaging has allowed better insights into the microarchitectural determinants of male bone strength. Specifically, androgens may promote trabecular bone development and thickness in young adulthood, as well as cortical consolidation in midlife and maintenance of cortical thickness and trabecular bone volume in older men by stimulating periosteal apposition and trabecular bone formation. Thus, although estrogen deficiency is the primary mediator of hypogonadal bone loss in men, high SHBG and low T probably pose additional detriments. The role of sex steroid catabolic enzymes and local lipid metabolism within bone (e.g. by 5 α -reductase type 1) requires further investigation. Recent studies using knockout mouse models have refined our understanding of the contributions of AR and ER α in osteoclasts, osteoblasts, and osteocytes to cortical




Figure 3: Schematic illustration of the cellular targets of androgen receptor (AR) and estrogen receptor (ER) α on periosteal, endocortical, and trabecular surfaces in male mice. Based on a μ CT image of the proximal murine tibia

and trabecular bone development and maintenance. At the same time, they have reinforced the notion that sex steroids must have important pleiotropic effects on bone, for example, via interaction with the nervous system, IGF-1 and altered response to mechanical loading. A better understanding of the microarchitectural, cellular and molecular mechanisms of actions of androgens and estrogens continues to be necessary to develop therapeutic strategies which can exploit their benefits for bone health without unwanted side effects in other tissues.

ACKNOWLEDGEMENTS

M. Laurent and V. Dubois are research fellows of the Research Foundation Flanders. This work was supported by the Research Foundation Flanders via grants G.0858.11N and G.0684.12N and OT/11/081 from KU Leuven.

COMPETING INTERESTS

Authors would declare that we have no financial or other competing interests.

REFERENCES

- Vanderschueren D, Vandenput L, Boonen S, Lindberg MK, Bouillon R, *et al*. Androgens and bone. *Endocr Rev* 2004; 25: 389–425.
- Laurent M, Gielen E, Claessens F, Boonen S, Vanderschueren D. Osteoporosis in older men: recent advances in pathophysiology and treatment. *Best Pract Res Clin Endocrinol Metab* 2013; 27: 527–39.
- Johnell O, Kanis JA. An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporos Int* 2006; 17: 1726–33.
- Kanis JA, Oden A, McCloskey EV, Johansson H, Wahl DA, *et al*. IOF Working Group on Epidemiology and Quality of Life. A systematic review of hip fracture incidence and probability of fracture worldwide. *Osteoporos Int* 2012; 23: 2239–56.
- Burge R, Dawson-Hughes B, Solomon DH, Wong JB, King A, *et al*. Incidence and economic burden of osteoporosis-related fractures in the United States, 2005–2025. *J Bone Miner Res* 2007; 22: 465–75.
- Tarride JE, Guo N, Hopkins R, Leslie WD, Morin S, *et al*. The burden of illness of osteoporosis in Canadian men. *J Bone Miner Res* 2012; 27: 1830–8.
- Tarride JE, Hopkins RB, Leslie WD, Morin S, Adachi JD, *et al*. The burden of illness of osteoporosis in Canada. *Osteoporos Int* 2012; 23: 2591–600.
- Frost M, Wraae K, Abrahamson B, Hoiberg M, Hagen C, *et al*. Osteoporosis and vertebral fractures in men aged 60–74 years. *Age Ageing* 2012; 41: 171–7.
- Nurmi-Luthje I, Sund R, Juntunen M, Luthje P. Post-hip fracture use of prescribed calcium plus vitamin D or vitamin D supplements and antiosteoporotic drugs is associated with lower mortality: a nationwide study in Finland. *J Bone Miner Res* 2011; 26: 1845–53.
- Wastesson JW, Ringback Weitoft G, Parker MG, Johnell K. educational level and use of osteoporosis drugs in elderly men and women: a Swedish nationwide register-based study. *Osteoporos Int* 2013; 24: 433–42.
- Curtis JR, McClure LA, Delzell E, Howard VJ, Orwoll E, *et al*. Population-based fracture risk assessment and osteoporosis treatment disparities by race and gender. *J Gen Intern Med* 2009; 24: 956–62.
- Shibli-Rahhal A, Vaughan-Sarrazin MS, Richardson K, Cram P. Testing and treatment for osteoporosis following hip fracture in an integrated U.S. healthcare delivery system. *Osteoporos Int* 2011; 22: 2973–80.
- Callewaert F, Boonen S, Vanderschueren D. Sex steroids and the male skeleton: a tale of two hormones. *Trends Endocrinol Metab* 2010; 21: 89–95.
- Sun L, Peng Y, Sharrow AC, Iqbal J, Zhang Z, *et al*. FSH directly regulates bone mass. *Cell* 2006; 125: 247–60.
- Zhu LL, Blair H, Cao J, Yuen T, Latif R, *et al*. Blocking antibody to the beta-subunit of FSH prevents bone loss by inhibiting bone resorption and stimulating bone synthesis. *Proc Natl Acad Sci U S A* 2012; 109: 14574–9.
- Yarram SJ, Perry MJ, Christopher TJ, Westby K, Brown NL, *et al*. Luteinizing hormone receptor knockout (Lurko) mice and transgenic human chorionic gonadotropin (Hcg)-overexpressing mice (Hcg Alpha-beta+) have bone phenotypes. *Endocrinology* 2003; 144: 3555–64.
- Seibel MJ, Dunstan CR, Zhou H, Allan CM, Handelsman DJ. Sex steroids, not FSH, influence bone mass. *Cell* 2006; 127: 1079.
- Allan CM, Kalak R, Dunstan CR, Mctavish KJ, Zhou H, *et al*. Follicle-stimulating hormone increases bone mass in female mice. *Proc Natl Acad Sci U S A* 2010; 107: 22629–34.
- Rouach V, Katzburg S, Koch Y, Stern N, Somjen D. Bone loss in ovariectomized rats: dominant role for estrogen but apparently not for FSH. *J Cell Biochem* 2011; 112: 128–37.
- Uihlein A, Kumbhani R, Siwila-Sackman E, Finkelstein J, Lee H, *et al*. FSH suppression in eugonadal men does not change bone turnover markers. *J Bone Miner Res* 2012; 27.
- Dunn JF, Nisula BC, Rodbard D. Transport of steroid hormones: binding of 21 endogenous steroids to both testosterone-binding globulin and corticosteroid-binding globulin in human plasma. *J Clin Endocrinol Metab* 1981; 53: 58–68.
- Mendel CM. The free hormone hypothesis: a physiologically based mathematical model. *Endocr Rev* 1989; 10: 232–74.
- Caldwell JD, Jirikowski GF. Sex hormone binding globulin and aging. *Horm Metab Res* 2009; 41: 173–82.
- Helsen C, Kerkhofs S, Clinckemalie L, Spans L, Laurent M, *et al*. Structural basis for nuclear hormone receptor DNA binding. *Mol Cell Endocrinol* 2012; 348: 411–7.
- Schauwaers K, De Gendt K, Saunders PT, Atanassova N, Haelens A, *et al*. Loss of androgen receptor binding to selective androgen response elements causes a reproductive phenotype in a knockin mouse model. *Proc Natl Acad Sci U S A* 2007; 104: 4961–6.
- Callewaert F, Sinnesael M, Gielen E, Boonen S, Vanderschueren D. Skeletal sexual dimorphism: relative contribution of sex steroids, GH-IGF1, and mechanical loading. *J Endocrinol* 2010; 207: 127–34.
- Ophoff J, Callewaert F, Venken K, De Gendt K, Ohlsson C, *et al*. Physical activity in the androgen receptor knockout mouse: evidence for reversal of androgen deficiency on cancellous bone. *Biochem Biophys Res Commun* 2009; 378: 139–44.
- Rana K, Fam BC, Clarke MV, Pang TP, Zajac JD, *et al*. Increased adiposity in dna binding-dependent androgen receptor knockout male mice associated with decreased voluntary activity and not insulin resistance. *Am J Physiol Endocrinol Metab* 2011; 301: E767–78.
- Boonen S, Dejaeger E, Vanderschueren D, Venken K, Bogaerts A, *et al*. Osteoporosis and osteoporotic fracture occurrence and prevention in the elderly: a geriatric perspective. *Best Pract Res Clin Endocrinol Metab* 2008; 22: 765–85.
- Masud T, Binkley N, Boonen S, Hannan MT. Official positions for frax (R) clinical regarding falls and frailty: can falls and frailty be used in frax (R)? from joint official positions development conference of the international society for clinical densitometry and international osteoporosis foundation on frax (R). *J Clin Densitom* 2011; 14: 194–204.
- Looker AC, Melton LJ 3rd, Harris T, Borrud L, Shepherd J, *et al*. Age, gender, and race/ethnic differences in total body and subregional bone density. *Osteoporos Int* 2009; 20: 1141–9.
- Meyer HE, Tverdal A, Falch JA. Risk factors for hip fracture in middle-aged norwegian women and men. *Am J Epidemiol* 1993; 137: 1203–11.
- Joakimsen RM, Fonnebo V, Magnus JH, Tollan A, Sogaard AJ. The tromsø study: body height, body mass index and fractures. *Osteoporos Int* 1998; 8: 436–42.
- Bjornerem A, Bui Q, Ghasem-Zadeh A, Hopper JL, Zebaze R, *et al*. Fracture risk and height: an association partly accounted for by cortical porosity of relatively thinner cortices. *J Bone Miner Res* 2013; 28: 2017–26.
- Rudang R, Mellstrom D, Clark E, Ohlsson C, Lorentzon M. Advancing maternal age is associated with lower bone mineral density in young adult male offspring. *Osteoporos Int* 2011; 23: 475–82.
- Cooper C, Westlake S, Harvey N, Javaid K, Dennison E, *et al*. Review: developmental origins of osteoporotic fracture. *Osteoporos Int* 2006; 17: 337–47.
- Hernandez CJ, Beaupre GS, Carter Dr. A theoretical analysis of the relative influences of peak BMD, age-related bone loss and menopause on the development of osteoporosis. *Osteoporos Int* 2003; 14: 843–7.
- Gafni RI, Baron J. Childhood bone mass acquisition and peak bone mass may not be important determinants of bone mass in late adulthood. *Pediatrics* 2007; 119 Suppl 2: S131–6.
- Baxter-Jones AD, Faulkner RA, Forwood MR, Mirwald RL, Bailey DA. Bone mineral accrual from 8 to 30 years of age: an estimation of peak bone mass. *J Bone Miner Res* 2011; 26: 1729–39.
- Ohlsson C, Darelid A, Nilsson M, Melin J, Mellstrom D, *et al*. Cortical consolidation due to increased mineralization and endosteal contraction in young adult men: a five-year longitudinal study. *J Clin Endocrinol Metab* 2011; 96: 2262–9.
- Berger C, Goltzman D, Langsetmo L, Joseph L, Jackson S, *et al*. CaMos Research Group. Peak bone mass from longitudinal data: implications for the prevalence, pathophysiology, and diagnosis of osteoporosis. *J Bone Miner Res* 2010; 25: 1948–57.
- Darelid A, Ohlsson C, Nilsson M, Kindblom JM, Mellstrom D, *et al*. Catch up in bone acquisition in young adult men with late normal puberty. *J Bone Miner Res* 2012; 27: 2198–207.
- Burghardt AJ, Kazakia GJ, Ramachandran S, Link TM, Majumdar S. Age- and gender-related differences in the geometric properties and biomechanical significance of intracortical porosity in the distal radius and tibia. *J Bone Miner Res* 2010; 25: 983–93.
- Macdonald HM, Nishiyama KK, Kang J, Hanley DA, Boyd SK. Age-related patterns of trabecular and cortical bone loss differ between sexes and skeletal sites: a population-based HR-pQCT study. *J Bone Miner Res* 2011; 26: 50–62.
- Szulc P, Munoz F, Duboeuf F, Marchand F, Delmas PD. Low width of tubular bones is associated with increased risk of fragility fracture in elderly men: the MINOS study. *Bone* 2006; 38: 595–602.
- Khosla S, Riggs BL, Atkinson EJ, Oberg AL, Mcdaniel LJ, *et al*. Effects of sex and age on bone microstructure at the ultradistal radius: a population-based noninvasive *in vivo* assessment. *J Bone Miner Res* 2006; 21: 124–31.

- 47 Dalzell N, Kaptoge S, Morris N, Berthier A, Koller B, *et al*. Bone micro-architecture and determinants of strength in the radius and tibia: age-related changes in a population-based study of normal adults measured with high-resolution pQCT. *Osteoporos Int* 2009; 20: 1683–94.
- 48 Kirmani S, Christen D, van Lenthe GH, Fischer PR, Bouxsein ML, *et al*. Bone structure at the distal radius during adolescent growth. *J Bone Miner Res* 2009; 24: 1033–42.
- 49 Walsh JS, Paggiosi MA, Eastell R. Cortical consolidation of the radius and tibia in young men and women. *J Clin Endocrinol Metab* 2012; 97: 3342–8.
- 50 Riggs BL, Melton LJ, Robb RA, Camp JJ, Atkinson EJ, *et al*. A population-based assessment of rates of bone loss at multiple skeletal sites: evidence for substantial trabecular bone loss in young adult women and men. *J Bone Miner Res* 2008; 23: 205–14.
- 51 Lauretani F, Bandinelli S, Griswold ME, Maggio M, Semba R, *et al*. Longitudinal changes in BMD and bone geometry in a population-based study. *J Bone Miner Res* 2008; 23: 400–8.
- 52 Duan Y, Beck TJ, Wang XF, Seeman E. Structural and biomechanical basis of sexual dimorphism in femoral neck fragility has its origins in growth and aging. *J Bone Miner Res* 2003; 18: 1766–74.
- 53 Riggs BL, Melton LJ 3rd, Robb RA, Camp JJ, Atkinson EJ, *et al*. Population-based study of age and sex differences in bone volumetric density, size, geometry, and structure at different skeletal sites. *J Bone Miner Res* 2004; 19: 1945–54.
- 54 Szulc P, Seeman E. Thinking inside and outside the envelopes of bone: dedicated to PDD. *Osteoporos Int* 2009; 20: 1281–8.
- 55 Szulc P, Boutroy S, Vilayphiou N, Chaitou A, Delmas PD, *et al*. Cross-sectional analysis of the association between fragility fractures and bone microarchitecture in older men: the STRAMBO study. *J Bone Miner Res* 2011; 26: 1358–67.
- 56 Peacock M, Buckwalter KA, Persohn S, Hangartner TN, Econs MJ, *et al*. Race and sex differences in bone mineral density and geometry at the femur. *Bone* 2009; 45: 218–25.
- 57 Srinivasan B, Kopperdahl DL, Amin S, Atkinson EJ, Camp J, *et al*. Relationship of femoral neck areal bone mineral density to volumetric bone mineral density, bone size, and femoral strength in men and women. *Osteoporos Int* 2011; 23: 155–62.
- 58 Riggs BL, Khosla S, Melton LJ 3rd. A unitary model for involutional osteoporosis: estrogen deficiency causes both type i and type ii osteoporosis in postmenopausal women and contributes to bone loss in aging men. *J Bone Miner Res* 1998; 13: 763–73.
- 59 Khosla S, Melton LJ 3rd, Riggs BL. The unitary model for estrogen deficiency and the pathogenesis of osteoporosis: is a revision needed? *J Bone Miner Res* 2011; 26: 441–51.
- 60 Manolagas SC. From estrogen-centric to aging and oxidative stress: a revised perspective of the pathogenesis of osteoporosis. *Endocr Rev* 2010; 31: 266–300.
- 61 Szulc P, Joly-Pharaboz MO, Marchand F, Delmas PD. Insulin-like growth factor i is a determinant of hip bone mineral density in men less than 60 years of age: MINOS study. *Calcif Tissue Int* 2004; 74: 322–9.
- 62 Ohlsson C, Mellstrom D, Carlzon D, Orwoll E, Ljunggren O, *et al*. Older men with low serum igf-1 have an increased risk of incident fractures: the MROS sweden study. *J Bone Miner Res* 2011; 26: 865–72.
- 63 Boonen S, Pye SR, O'Neill TW, Szulc P, Gielen E, *et al*. EMAS Group. Influence of bone remodelling rate on quantitative ultrasound parameters at the calcaneus and dxa bmda of the hip and spine in middle-aged and elderly european men: The European Male Ageing Study (EMAS). *Eur J Endocrinol* 2011; 165: 977–86.
- 64 Pye SR, Almusalam B, Boonen S, Vanderschueren D, Borghs H, *et al*. EMAS Study Group. Influence of insulin-like growth factor binding protein (igfbp)-1 and igfbp-3 on bone health: results from the European Male Ageing Study. *Calcif Tissue Int* 2011; 88: 503–10.
- 65 Curtis JR, Ewing SK, Bauer DC, Cauley JA, Cawthon PM, *et al*. Association of intact parathyroid hormone levels with subsequent hip BMD loss: The Osteoporotic Fractures in men (MROS) study. *J Clin Endocrinol Metab* 2012; 97: 1937–44.
- 66 Grundberg E, Lau EM, Pastinen T, Kindmark A, Nilsson O, *et al*. Vitamin d receptor 3' haplotypes are unequally expressed in primary human bone cells and associated with increased fracture risk: The MROS study in Sweden and Hong Kong. *J Bone Miner Res* 2007; 22: 832–40.
- 67 Vanderschueren D, Pye SR, O'Neill TW, Lee DM, Jans I, *et al*. EMAS Study Group. Active vitamin d (1,25-dihydroxyvitamin d) and bone health in middle-aged and elderly men: The European Male Ageing Study (EMAS). *J Clin Endocrinol Metab* 2013; 98: 995–1005.
- 68 Roef G, Lapauw B, Goemaere S, Zmierzczak H, Fiers T, *et al*. thyroid hormone status within the physiological range affects bone mass and density in healthy men at the age of peak bone mass. *Eur J Endocrinol* 2011; 164: 1027–34.
- 69 Waring AC, Harrison S, Fink HA, Samuels MH, Cawthon PM, *et al*. Osteoporotic Fractures in Men (MROS) Study. A prospective study of thyroid function, bone loss, and fractures in older men: The MROS study. *J Bone Miner Res* 2013; 28: 472–9.
- 70 Rolland T, Boutroy S, Vilayphiou N, Blaizot S, Chapurlat R, *et al*. Poor trabecular microarchitecture at the distal radius in older men with increased concentration of high-sensitivity c-reactive protein: The STRAMBO study. *Calcif Tissue Int* 2012; 90: 496–506.
- 71 Roshandel D, Holliday KL, Pye SR, Boonen S, Borghs H, *et al*. EMAS Study Group. Genetic variation in the rankl/rank/opg signaling pathway is associated with bone turnover and bone mineral density in men. *J Bone Miner Res* 2010; 25: 1830–8.
- 72 Roshandel D, Holliday KL, Pye SR, Ward KA, Boonen S, *et al*. EMAS Study Group. Influence of polymorphisms in the RANKL/RANK/OPG signaling pathway on volumetric bone mineral density and bone geometry at the forearm in men. *Calcif Tissue Int* 2011; 89: 446–55.
- 73 Roshandel D, Thomson W, Pye SR, Boonen S, Borghs H, *et al*. EMAS Study Group. Polymorphisms in genes involved in the NF- κ B signalling pathway are associated with bone mineral density, geometry and turnover in men. *PLoS One* 2011; 6: e28031.
- 74 Liu CT, Estrada K, Yerges-Armstrong LM, Amin N, Evangelou E, *et al*. Assessment of gene-by-sex interaction effect on bone mineral density. *J Bone Miner Res* 2012; 27: 2051–64.
- 75 Van Caenegem E, Wierckx K, Taes Y, Dedecker D, Van De Peer F, *et al*. Bone mass, bone geometry, and body composition in female-to-male transsexual persons after long-term cross-sex hormonal therapy. *J Clin Endocrinol Metab* 2012; 97: 2503–11.
- 76 Taes Y, Lapauw B, Vandewalle S, Zmierzczak H, Goemaere S, *et al*. Estrogen-specific action on bone geometry and volumetric bone density: longitudinal observations in an adult with complete androgen insensitivity. *Bone* 2009; 45: 392–7.
- 77 Han TS, Goswami D, Trikudanathan S, Creighton SM, Conway GS. Comparison of bone mineral density and body proportions between women with complete androgen insensitivity syndrome and women with gonadal dysgenesis. *Eur J Endocrinol* 2008; 159: 179–85.
- 78 Almeida M, Han L, Martin-Millan M, Plotkin LI, Stewart SA, *et al*. Skeletal involution by age-associated oxidative stress and its acceleration by loss of sex steroids. *J Biol Chem* 2007; 282: 27285–97.
- 79 Bellido T, Jilka RL, Boyce BF, Girasole G, Broxmeyer H, *et al*. Regulation of interleukin-6, osteoclastogenesis, and bone mass by androgens. The role of the androgen receptor. *J Clin Invest* 1995; 95: 2886–95.
- 80 Hofbauer LC, Ten RM, Khosla S. The anti-androgen hydroxyflutamide and androgens inhibit interleukin-6 production by an androgen-responsive human osteoblastic cell line. *J Bone Miner Res* 1999; 14: 1330–7.
- 81 Callewaert F, Venken K, Kopchick JJ, Torcaso A, van Lenthe GH, *et al*. Sexual dimorphism in cortical bone size and strength but not density is determined by independent and time-specific actions of sex steroids and IGF-1: evidence from pubertal mouse models. *J Bone Miner Res* 2010; 25: 617–26.
- 82 Perry RJ, Farquharson C, Ahmed SF. The role of sex steroids in controlling pubertal growth. *Clin Endocrinol (Oxf)* 2008; 68: 4–15.
- 83 Smith EP, Boyd J, Frank GR, Takahashi H, Cohen RM, *et al*. Estrogen resistance caused by a mutation in the estrogen-receptor gene in a man. *N Engl J Med* 1994; 331: 1056–61.
- 84 Carani C, Qin K, Simoni M, Faustini-Fustini M, Serpente S, *et al*. Effect of testosterone and estradiol in a man with aromatase deficiency. *N Engl J Med* 1997; 337: 91–5.
- 85 Bilezikian JP, Morishima A, Bell J, Grumbach MM. Increased bone mass as a result of estrogen therapy in a man with aromatase deficiency. *N Engl J Med* 1998; 339: 599–603.
- 86 Bouillon R, Bex M, Vanderschueren D, Boonen S. Estrogens are essential for male pubertal periosteal bone expansion. *J Clin Endocrinol Metab* 2004; 89: 6025–9.
- 87 Merlotti D, Gennari L, Stolakis K, Nuti R. Aromatase activity and bone loss in men. *J Osteoporos* 2011; 2011: 230671.
- 88 Borjesson AE, Windahl SH, Karimian E, Eriksson EE, Lagerquist MK, *et al*. The role of estrogen receptor-alpha and its activation function-1 for growth plate closure in female mice. *Am J Physiol Endocrinol Metab* 2012; 302: E1381–9.
- 89 Zachmann M, Prader A, Sobel EH, Crigler JF Jr, Ritzen EM, *et al*. pubertal growth in patients with androgen insensitivity: indirect evidence for the importance of estrogens in pubertal growth of girls. *J Pediatr* 1986; 108: 694–7.
- 90 Callewaert F, Venken K, Ophoff J, De Gendt K, Torcaso A, *et al*. Differential regulation of bone and body composition in male mice with combined inactivation of androgen and estrogen receptor-alpha. *FASEB J* 2009; 23: 232–40.
- 91 Albright F, Smith PH, Richardson AM. Postmenopausal osteoporosis. *JAMA* 1941; 116: 2465–74.
- 92 Finkelstein JS, Klibanski A, Neer RM, Greenspan SL, Rosenthal DI, *et al*. Osteoporosis in men with idiopathic hypogonadotropic hypogonadism. *Ann Intern Med* 1987; 106: 354–61.
- 93 Swartz CM, Young MA. Male hypogonadism and bone fracture. *N Engl J Med* 1988; 318: 996.
- 94 Slemenda CW, Longcope C, Zhou L, Hui SL, Peacock M, *et al*. Sex steroids and bone mass in older men. Positive Associations with serum estrogens and negative associations with androgens. *J Clin Invest* 1997; 100: 1755–9.
- 95 Khosla S, Melton LJ 3rd, Atkinson EJ, O'fallon WM, Klee GG, *et al*. Relationship of serum sex steroid levels and bone turnover markers with bone mineral density in men and women: a key role for bioavailable estrogen. *J Clin Endocrinol Metab* 1998; 83: 2266–74.
- 96 Khosla S, Melton LJ 3rd, Atkinson EJ, O'fallon WM. Relationship of serum sex steroid levels to longitudinal changes in bone density in young versus elderly men. *J Clin Endocrinol Metab* 2001; 86: 3555–61.
- 97 Szulc P, Munoz F, Claustrat B, Garnero P, Marchand F, *et al*. Bioavailable estradiol

- may be an important determinant of osteoporosis in men: the MINOS study. *J Clin Endocrinol Metab* 2001; 86: 192–9.
- 98 Gennari L, Merlotti D, Martini G, Gonnelli S, Franci B, *et al*. Longitudinal association between sex hormone levels, bone loss, and bone turnover in elderly men. *J Clin Endocrinol Metab* 2003; 88: 5327–33.
 - 99 Szulc P, Uusi-Rasi K, Claustrat B, Marchand F, Beck TJ, *et al*. Role of sex steroids in the regulation of bone morphology in men. The MINOS study. *Osteoporos Int* 2004; 15: 909–17.
 - 100 Goderie-Plomp HW, van der Klift M, De Ronde W, Hofman A, de jong FH, *et al*. Endogenous sex hormones, sex hormone-binding globulin, and the risk of incident vertebral fractures in elderly men and women: The Rotterdam Study. *J Clin Endocrinol Metab* 2004; 89: 3261–9.
 - 101 Khosla S, Melton LJ 3rd, Robb RA, Camp JJ, Atkinson EJ, *et al*. Relationship of volumetric bmd and structural parameters at different skeletal sites to sex steroid levels in men. *J Bone Miner Res* 2005; 20: 730–40.
 - 102 Vandenput L, Labrie F, Mellstrom D, Swanson C, Knutsson T, *et al*. Serum levels of specific glucuronidated androgen metabolites predict bmd and prostate volume in elderly men. *J Bone Miner Res* 2007; 22: 220–7.
 - 103 Kuchuk NO, van Schoor NM, Pluijm SM, Smit JH, de Ronde W, *et al*. The association of sex hormone levels with quantitative ultrasound, bone mineral density, bone turnover and osteoporotic fractures in older men and women. *Clin Endocrinol (Oxf)* 2007; 67: 295–303.
 - 104 Araujo AB, Travison TG, Leder BZ, Mckinlay JB. Correlations between serum testosterone, estradiol, and sex hormone-binding globulin and bone mineral density in a diverse sample of men. *J Clin Endocrinol Metab* 2008; 93: 2135–41.
 - 105 Vanderschueren D, Pye SR, Venken K, Borghs H, Gaytant J, *et al*. EMAS Study Group. Gonadal sex steroid status and bone health in middle-aged and elderly European men. *Osteoporos Int* 2010; 21: 1331–9.
 - 106 Mellstrom D, Vandenput L, Mallmin H, Holmberg AH, Lorentzon M, *et al*. Older men with low serum estradiol and high serum shbg have an increased risk of fractures. *J Bone Miner Res* 2008; 23: 1552–60.
 - 107 Ward KA, Pye SR, Adams JE, Boonen S, Vanderschueren D, *et al*. EMAS Study Group. Influence of age and sex steroids on bone density and geometry in middle-aged and elderly European men. *Osteoporos Int* 2011; 22: 1513–23.
 - 108 Wu FC, Tajar A, Beynon JM, Pye SR, Silman AJ, *et al*. EMAS Group. Identification of late-onset hypogonadism in middle-aged and elderly men. *N Engl J Med* 2010; 363: 123–35.
 - 109 Vandenput L, Boonen S, Van Herck E, Swinnen JV, Bouillon R, *et al*. Evidence from the aged orchidectomized male rat model that 17beta-estradiol is a more effective bone-sparing and anabolic agent than 5alpha-dihydrotestosterone. *J Bone Miner Res* 2002; 17: 2080–6.
 - 110 Limer KL, Pye SR, Thomson W, Boonen S, Borghs H, *et al*. EMAS Study Group. Genetic variation in sex hormone genes influences heel ultrasound parameters in middle-aged and elderly men: results from the European Male Aging Study (EMAS). *J Bone Miner Res* 2009; 24: 314–23.
 - 111 Holliday KL, Pye SR, Thomson W, Boonen S, Borghs H, *et al*. EMAS Study Group. The ESR1 (6q25) locus is associated with calcaneal ultrasound parameters and radial volumetric bone mineral density in European men. *Plos One* 2011; 6: e22037.
 - 112 Huhtaniemi IT, Pye SR, Holliday KL, Thomson W, O'Neill TW, *et al*. European Male Aging Study Group. Effect of polymorphisms in selected genes involved in pituitary-testicular function on reproductive hormones and phenotype in aging men. *J Clin Endocrinol Metab* 2010; 95: 1898–908.
 - 113 Huhtaniemi IT, Pye SR, Limer KL, Thomson W, O'Neill TW, *et al*. European Male Aging Study Group. Increased estrogen rather than decreased androgen action is associated with longer androgen receptor CAG repeats. *J Clin Endocrinol Metab* 2009; 94: 277–84.
 - 114 Shahinian VB, Kuo YF, Freeman JL, Goodwin JS. Risk of fracture after androgen deprivation for prostate cancer. *N Engl J Med* 2005; 352: 154–64.
 - 115 Smith MR, Goode M, Zietman AL, MCGovern FJ, Lee H, *et al*. Bicalutamide monotherapy versus leuprolide monotherapy for prostate cancer: effects on bone mineral density and body composition. *J Clin Oncol* 2004; 22: 2546–53.
 - 116 Sieber PR, Keiller DL, Kahnoski RJ, Gallo J, McFadden S. Bicalutamide 150 mg maintains bone mineral density during monotherapy for localized or locally advanced prostate cancer. *J Urol* 2004; 171: 2272–6.
 - 117 Wadhwa VK, Weston R, Mistry R, Parr NJ. Long-term changes in bone mineral density and predicted fracture risk in patients receiving androgen-deprivation therapy for prostate cancer, with stratification of treatment based on presenting values. *BJU Int* 2009; 104: 800–5.
 - 118 Burnett-Bowie SA, Mckay EA, Lee H, Leder BZ. Effects of aromatase inhibition on bone mineral density and bone turnover in older men with low testosterone levels. *J Clin Endocrinol Metab* 2009; 94: 4785–92.
 - 119 Falahati-Nini A, Riggs BL, Atkinson EJ, O'Fallon WM, Eastell R, *et al*. Relative contributions of testosterone and estrogen in regulating bone resorption and formation in normal elderly men. *J Clin Invest* 2000; 106: 1553–60.
 - 120 Yu E, Taylor A, Wulczyn K, Webb M, Perros N, *et al*. Hypogonadism with estrogen removal (Her): differential effects of androgens and estrogens on bone microarchitecture in adult men. *J Bone Mineral Res* 2012; 27: S66.
 - 121 Vandenput L, Ederveen AG, Erben RG, Stahr K, Swinnen JV, *et al*. Testosterone prevents orchidectomy-induced bone loss in estrogen receptor-alpha knockout mice. *Biochem Biophys Res Commun* 2001; 285: 70–6.
 - 122 Mellstrom D, Johnell O, Ljunggren O, Eriksson AL, Lorentzon M, *et al*. Free testosterone is an independent predictor of BMD and prevalent fractures in elderly men: MROS Sweden. *J Bone Miner Res* 2006; 21: 529–35.
 - 123 Meier C, Nguyen TV, Handelsman DJ, Schindler C, Kushnir MM, *et al*. Endogenous sex hormones and incident fracture risk in older men: The Dubbo Osteoporosis Epidemiology Study. *Arch Intern Med* 2008; 168: 47–54.
 - 124 van den Beld AW, de Jong FH, Grobbee DE, Pols HA, Lamberts SW. Measures of bioavailable serum testosterone and estradiol and their relationships with muscle strength, bone density, and body composition in elderly men. *J Clin Endocrinol Metab* 2000; 85: 3276–82.
 - 125 Tajar A, Huhtaniemi IT, O'Neill TW, Finn JD, Pye SR, *et al*. EMAS Group. Characteristics of androgen deficiency in late-onset hypogonadism: results from the European male aging study (EMAS). *J Clin Endocrinol Metab* 2012; 97: 1508–16.
 - 126 Amin S, Zhang Y, Felson DT, Sawin CT, Hannan MT, *et al*. Estradiol, testosterone, and the risk for hip fractures in elderly men from the framingham study. *Am J Med* 2006; 119: 426–33.
 - 127 Vandenput L, Swinnen JV, Boonen S, Van Herck E, Erben RG, *et al*. Role of the androgen receptor in skeletal homeostasis: the androgen-resistant testicular feminized male mouse model. *J Bone Miner Res* 2004; 19: 1462–70.
 - 128 Rochira V, Zirilli L, Madeo B, Aranda C, Caffagni G, *et al*. skeletal effects of long-term estrogen and testosterone replacement treatment in a man with congenital aromatase deficiency: evidences of a priming effect of estrogen for sex steroids action on bone. *Bone* 2007; 40: 1662–8.
 - 129 Leblanc ES, Nielson CM, Marshall LM, Lapidus JA, Barrett-Connor E, *et al*. Osteoporotic Fractures in Men Study Group. The effects of serum testosterone, estradiol, and sex hormone binding globulin levels on fracture risk in older men. *J Clin Endocrinol Metab* 2009; 94: 3337–46.
 - 130 Cauley JA, Ewing SK, Taylor BC, Fink HA, Ensrud KE, *et al*. Osteoporotic Fractures in Men Study (MrOS) Research Group. Sex steroid hormones in older men: longitudinal associations with 4.5-year change in hip bone mineral density—the osteoporotic fractures in men study. *J Clin Endocrinol Metab* 2010; 95: 4314–23.
 - 131 Vanbillemont G, Lapauw B, Bogaert V, Goemaere S, Zmierzczak HG, *et al*. Sex hormone-binding globulin as an independent determinant of cortical bone status in men at the age of peak bone mass. *J Clin Endocrinol Metab* 2010; 95: 1579–86.
 - 132 Lorentzon M, Swanson C, Andersson N, Mellstrom D, Ohlsson C. Free testosterone is a positive, whereas free estradiol is a negative, predictor of cortical bone size in young Swedish men: The GOOD study. *J Bone Miner Res* 2005; 20: 1334–41.
 - 133 Eriksson AL, Lorentzon M, Mellstrom D, Vandenput L, Swanson C, *et al*. SHBG gene promoter polymorphisms in men are associated with serum sex hormone-binding globulin, androgen and androgen metabolite levels, and hip bone mineral density. *J Clin Endocrinol Metab* 2006; 91: 5029–37.
 - 134 Center JR, Nguyen TV, Sambrook PN, Eisman JA. Hormonal and biochemical parameters and osteoporotic fractures in elderly men. *J Bone Miner Res* 2000; 15: 1405–11.
 - 135 Khosla S. Editorial: sex hormone binding globulin: inhibitor or facilitator (or both) of sex steroid action? *J Clin Endocrinol Metab* 2006; 91: 4764–6.
 - 136 Swanson C, Lorentzon M, Vandenput L, Labrie F, Rane A, *et al*. Sex steroid levels and cortical bone size in young men are associated with a uridine diphosphate glucuronosyltransferase 2B7 polymorphism (H268Y). *J Clin Endocrinol Metab* 2007; 92: 3697–704.
 - 137 Lorentzon M, Eriksson AL, Mellstrom D, Ohlsson C. The COMT Val158met polymorphism is associated with peak BMD in men. *J Bone Miner Res* 2004; 19: 2005–11.
 - 138 Eriksson AL, Mellstrom D, Lorentzon M, Orwoll ES, Redlund-Johnell I, *et al*. The COMT Val158met polymorphism is associated with prevalent fractures in Swedish men. *Bone* 2008; 42: 107–12.
 - 139 Lorentzon M, Eriksson AL, Nilsson S, Mellstrom D, Ohlsson C. Association between physical activity and BMD in young men is modulated by catechol-o-methyltransferase (COMT) genotype: The GOOD study. *J Bone Miner Res* 2007; 22: 1165–72.
 - 140 Stolck L, van Meurs JB, Jhamai M, Arp PP, van Leeuwen JP, *et al*. The catechol-o-methyltransferase met158 low-activity allele and association with nonvertebral fracture risk in elderly men. *J Clin Endocrinol Metab* 2007; 92: 3206–12.
 - 141 Issa S, Schnabel D, Feix M, Wolf L, Schaefer HE, *et al*. Human osteoblast-like cells express predominantly steroid 5alpha-reductase type 1. *J Clin Endocrinol Metab* 2002; 87: 5401–7.
 - 142 Tollin SR, Rosen HN, Zurowski K, Saltzman B, Zeind AJ, *et al*. Finasteride therapy does not alter bone turnover in men with benign prostatic hyperplasia—a Clinical Research Center Study. *J Clin Endocrinol Metab* 1996; 81: 1031–4.
 - 143 Matsumoto AM, Tenover L, Mcclung M, Mobley D, Geller J, *et al*. Pless Study Group. The long-term effect of specific type ii 5alpha-reductase inhibition with finasteride on bone mineral density in men: results of a 4-year placebo controlled trial. *J Urol* 2002; 167: 2105–8.
 - 144 Amory JK, Watts NB, Easley KA, Sutton PR, Anawalt BD, *et al*. Exogenous

- testosterone or testosterone with finasteride increases bone mineral density in older men with low serum testosterone. *J Clin Endocrinol Metab* 2004; 89: 503–10.
- 145 Amory JK, Anawalt BD, Matsumoto AM, Page ST, Bremner WJ, *et al*. The effect of 5 α -reductase inhibition with dutasteride and finasteride on bone mineral density, serum lipoproteins, hemoglobin, prostate specific antigen and sexual function in healthy young men. *J Urol* 2008; 179: 2333–8.
- 146 Jacobsen SJ, Cheetham TC, Haque R, Shi JM, Loo RK. Association between 5 α -reductase inhibition and risk of hip fracture. *JAMA* 2008; 300: 1660–4.
- 147 Vestergaard P, Rejnmark L, Mosekilde L. Risk of fractures associated with treatment for benign prostate hyperplasia in men. *Osteoporos Int* 2011; 22: 731–7.
- 148 Page ST, Amory JK, Bowman FD, Anawalt BD, Matsumoto AM, *et al*. Exogenous testosterone (t) alone or with finasteride increases physical performance, grip strength, and lean body mass in older men with low serum T. *J Clin Endocrinol Metab* 2005; 90: 1502–10.
- 149 Windahl SH, Andersson N, Borjesson AE, Swanson C, Svensson J, *et al*. Reduced bone mass and muscle strength in male 5 α -reductase type 1 inactivated mice. *PLoS One* 2011; 6: e21402.
- 150 Andriole GL, Kirby R. Safety and tolerability of the dual 5 α -reductase inhibitor dutasteride in the treatment of benign prostatic hyperplasia. *Eur Urol* 2003; 44: 82–8.
- 151 Bhasin S, Travison TG, Storer TW, Lakshman K, Kaushik M, *et al*. Effect of testosterone supplementation with and without a dual 5 α -reductase inhibitor on fat-free mass in men with suppressed testosterone production: a randomized controlled trial. *JAMA* 2012; 307: 931–9.
- 152 Meier C, Liu PY, Ly LP, De Winter-Modzelewski J, Jimenez M, *et al*. Recombinant human chorionic gonadotropin but not dihydrotestosterone alone stimulates osteoblastic collagen synthesis in older men with partial age-related androgen deficiency. *J Clin Endocrinol Metab* 2004; 89: 3033–41.
- 153 Idan A, Griffiths KA, Harwood DT, Seibel MJ, Turner L, *et al*. Long-term effects of dihydrotestosterone treatment on prostate growth in healthy, middle-aged men without prostate disease: a randomized, placebo-controlled trial. *Ann Intern Med* 2010; 153: 621–32.
- 154 Moverare S, Venken K, Eriksson AL, Andersson N, Skrtic S, *et al*. Differential effects on bone of estrogen receptor alpha and androgen receptor activation in orchidectomized adult male mice. *Proc Natl Acad Sci U S A* 2003; 100: 13573–8.
- 155 Tivesten A, Moverare-Skrtic S, Chagin A, Venken K, Salmon P, *et al*. Additive protective effects of estrogen and androgen treatment on trabecular bone in ovariectomized rats. *J Bone Miner Res* 2004; 19: 1833–9.
- 156 Venken K, De Gendt K, Boonen S, Ophoff J, Bouillon R, *et al*. Relative impact of androgen and estrogen receptor activation in the effects of androgens on trabecular and cortical bone in growing male mice: a study in the androgen receptor knockout mouse model. *J Bone Miner Res* 2006; 21: 576–85.
- 157 Sims NA, Dupont S, Krust A, Clement-Lacroix P, Minet D, *et al*. Deletion of estrogen receptors reveals a regulatory role for estrogen receptors-beta in bone remodeling in females but not in males. *Bone* 2002; 30: 18–25.
- 158 Maclean HE, Moore AJ, Sastra SA, Morris HA, Ghasem-Zadeh A, *et al*. DNA-binding-dependent androgen receptor signaling contributes to gender differences and has physiological actions in males and females. *J Endocrinol* 2010; 206: 93–103.
- 159 Windahl SH, Vidal O, Andersson G, Gustafsson JA, Ohlsson C. Increased cortical bone mineral content but unchanged trabecular bone mineral density in female erbeta(-/-) mice. *J Clin Invest* 1999; 104: 895–901.
- 160 Sims NA, Clement-Lacroix P, Minet D, Fraslon-Vanhulle C, Gaillard-Kelly M, *et al*. A functional androgen receptor is not sufficient to allow estradiol to protect bone after gonadectomy in estradiol receptor-deficient mice. *J Clin Invest* 2003; 111: 1319–27.
- 161 Kawano H, Sato T, Yamada T, Matsumoto T, Sekine K, *et al*. Suppressive function of androgen receptor in bone resorption. *Proc Natl Acad Sci U S A* 2003; 100: 9416–21.
- 162 Chiang C, Chiu M, Moore AJ, Anderson PH, Ghasem-Zadeh A, *et al*. Mineralization and bone resorption are regulated by the androgen receptor in male mice. *J Bone Miner Res* 2009; 24: 621–31.
- 163 Notini AJ, Mcmanus JF, Moore A, Bouxsein M, Jimenez M, *et al*. Osteoblast deletion of exon 3 of the androgen receptor gene results in trabecular bone loss in adult male mice. *J Bone Miner Res* 2007; 22: 347–56.
- 164 Wiren KM, Zhang XW, Toombs AR, Kasparcova V, Gentile MA, *et al*. Targeted overexpression of androgen receptor in osteoblasts: unexpected complex bone phenotype in growing animals. *Endocrinology* 2004; 145: 3507–22.
- 165 Wiren KM, Semirale AA, Zhang XW, Woo A, Tommasini SM, *et al*. Targeting of androgen receptor in bone reveals a lack of androgen anabolic action and inhibition of osteogenesis: a model for compartment-specific androgen action in the skeleton. *Bone* 2008; 43: 440–51.
- 166 Sinnesael M, Claessens F, Laurent M, Dubois V, Boonen S, *et al*. Androgen receptor (AR) in osteocytes is important for the maintenance of male skeletal integrity: evidence from targeted AR disruption in mouse osteocytes. *J Bone Miner Res* 2012; 27: 2535–43.
- 167 Abu EO, Horner A, Kusec V, Triffitt JT, Compston JE. The localization of androgen receptors in human bone. *J Clin Endocrinol Metab* 1997; 82: 3493–7.
- 168 Turner AG, Notini AJ, Chiu WS, Hodge J, Nicholson G, *et al*. Androgen receptor expression and function in osteoclasts. *Open Physiol J* 2008; 1.
- 169 Huber DM, Bendixen AC, Pathrose P, Srivastava S, Dienger KM, *et al*. Androgens suppress osteoclast formation induced by rankl and macrophage-colony stimulating factor. *Endocrinology* 2001; 142: 3800–8.
- 170 Pederson L, Kremer M, Judd J, Pascoe D, Spelsberg TC, *et al*. Androgens regulate bone resorption activity of isolated osteoclasts *in vitro*. *Proc Natl Acad Sci U S A* 1999; 96: 505–10.
- 171 Michael H, Harkonen PL, Vaananen HK, Hentunen TA. Estrogen and testosterone use different cellular pathways to inhibit osteoclastogenesis and bone resorption. *J Bone Miner Res* 2005; 20: 2224–32.
- 172 Nakamura T, Watanabe T, Nakamichi Y, Fukuda T, Matsumoto T, *et al*. Genetic evidence of androgen receptor function in osteoclasts: generation and characterization of osteoclast-specific androgen receptor knockout mice. *J Bone Miner Res* 2004; 19: S3.
- 173 Nakamura T, Imai Y, Matsumoto T, Sato S, Takeuchi K, *et al*. Estrogen prevents bone loss via estrogen receptor alpha and induction of fas ligand in osteoclasts. *Cell* 2007; 130: 811–23.
- 174 Martin-Millan M, Almeida M, Ambrogini E, Han L, Zhao H, *et al*. The estrogen receptor-alpha in osteoclasts mediates the protective effects of estrogens on cancellous but not cortical bone. *Mol Endocrinol* 2010; 24: 323–34.
- 175 Sjogren K, Lagerquist M, Moverare-Skrtic S, Andersson N, Windahl SH, *et al*. Elevated aromatase expression in osteoblasts leads to increased bone mass without systemic adverse effects. *J Bone Miner Res* 2009; 24: 1263–70.
- 176 Almeida M, Iyer S, Martin-Millan M, Bartell SM, Han L, *et al*. Estrogen receptor-alpha signaling in osteoblast progenitors stimulates cortical bone accrual. *J Clin Invest* 2013; 123: 394–404.
- 177 Maatta JA, Buki KG, Gu G, Alanne MH, Vaaranen J, *et al*. Inactivation of estrogen receptor alpha in bone-forming cells induces bone loss in female mice. *FASEB J* 2010; 24: 478–88.
- 178 Windahl SH, Borjesson AE, Farman HH, Engdahl C, Moverare-Skrtic S, *et al*. Estrogen receptor-alpha in osteocytes is important for trabecular bone formation in male mice. *Proc Natl Acad Sci U S A* 2013; 110: 2294–9.
- 179 Lee K, Jessop H, Suswillo R, Zaman G, Lanyon L. Endocrinology: bone adaptation requires oestrogen receptor-alpha. *Nature* 2003; 424: 389.
- 180 Callewaert F, Bakker A, Schrooten J, Van Meerbeek B, Verhoeven G, *et al*. Androgen receptor disruption increases the osteogenic response to mechanical loading in male mice. *J Bone Miner Res* 2010; 25: 124–31.
- 181 MacLean HE, Chiu WS, Notini AJ, Axell AM, Davey RA, *et al*. Impaired skeletal muscle development and function in male, but not female, genomic androgen receptor knockout mice. *FASEB J* 2008; 22: 2676–89.
- 182 Ophoff J, Van Proeyen K, Callewaert F, De Gendt K, De Bock K, *et al*. Androgen signaling in myocytes contributes to the maintenance of muscle mass and fiber type regulation but not to muscle strength or fatigue. *Endocrinology* 2009; 150: 3558–66.
- 183 Chambon C, Duteil D, Vignaud A, Ferry A, Messaddeq N, *et al*. Myocytic androgen receptor controls the strength but not the mass of limb muscles. *Proc Natl Acad Sci U S A* 2010; 107: 14327–32.
- 184 Dubois V, Laurent M, Boonen S, Vanderschueren D, Claessens F. Androgens and skeletal muscle: cellular and molecular action mechanisms underlying the anabolic actions. *Cell Mol Life Sci* 2012; 69: 1651–67.
- 185 Gielen E, Verschueren S, O'Neill TW, Pye SR, O'Connell MD, *et al*. Musculoskeletal frailty: a geriatric syndrome at the core of fracture occurrence in older age. *Calcif Tissue Int* 2012; 91: 161–77.
- 186 Verschueren S, Gielen E, O'Neill TW, Pye SR, Adams JE, *et al*. Sarcopenia and its relationship with bone mineral density in middle-aged and elderly European men. *Osteoporos Int* 2013; 24: 87–98.
- 187 Ohlsson C, Engdahl C, Borjesson AE, Windahl SH, Studer E, *et al*. Estrogen receptor-alpha expression in neuronal cells affects bone mass. *Proc Natl Acad Sci U S A* 2012; 109: 983–8.
- 188 Raskin K, de Gendt K, Duittoz A, Liere P, Verhoeven G, *et al*. Conditional inactivation of androgen receptor gene in the nervous system: effects on male behavioral and neuroendocrine responses. *J Neurosci* 2009; 29: 4461–70.
- 189 Olson LE, Ohlsson C, Mohan S. The role of GH/IGF-I-mediated mechanisms in sex differences in cortical bone size in mice. *Calcif Tissue Int* 2011; 88: 1–8.

How to cite this article: Laurent M, Antonio L, Sinnesael M, Dubois V, Gielen E, Claessens F, Vanderschueren D. Androgens and estrogens in skeletal sexual dimorphism. *Asian J Androl* 20 January 2014. doi: 10.4103/1008-682X.122356. [Epub ahead of print]

